Abbreviations

ABPM ambulatory blood pressure monitoring ACE inhibitor angiotensin converting enzyme inhibitor

ACOG American College of Obstetricians and Gynecologists

ACR albumin : creatinine ratio
AFI amniotic fluid index

AFLP acute fatty liver of pregnancy

ANC antenatal care

AOM Association of Ontario Midwives APTT activated partial thromboplastin time

ARB angiotensin receptor blocker AST aspartate transaminase

BEmONC basic emergency obstetric and neonatal care

BID twice daily dosing
BMI body mass index
BP blood pressure
BPP biophysical profile
CBC complete blood count

CEmONC comprehensive emergency obstetric and neonatal care

cGMP cyclic guanosine monophosphate

CO carbon monoxide CPG clinical practice guideline

CSE combined spinal-epidural analgesia

CVP central venous pressure

DASH dietary approaches to stop hypertension

dBP diastolic blood pressure

DIC disseminated intravascular coagulation
EmONC emergency obstetric and neonatal care
eNOS endothelial nitric oxide synthase
GDM gestational diabetes mellitus
GH gestational hypertension
GMP guanosine monophosphate

GRADE grades of recommendation, assessment, development and evaluation

GSNO S-nitrosoglutathione
GTP guanosine triphosphate
HDL high-density lipoprotein

HDP hypertensive disorder of pregnancy

HELLP syndrome haemolysis, elevated liver enzymes and low platelets syndrome

HIC high-income country

HR heart rate IM intramuscular

INR international normalised ratio

IOL induction of labour

ISSHP International Society for the Study of Hypertension in Pregnancy

IUGR intrauterine fetal growth restriction

IV intravenous

LDH lactate dehydrogenase LDL low-density liporotein

LMICs low- and middle-income countries

LMP last menstrual period

MEOWS modified early obstetric warning systems

MRI magnetic resonance imaging

NICE National Institute for Health and Clinical Excellence

NICU neonatal intensive care unit

NO nitric oxide

NSAIDs non-steroidal anti-inflammatory drugs

NST non-stress test

NVOG National Obstetrics and Gynaecology Society, The Netherlands

PCA patient-controlled labour analgesia

PDE5 phosphodiesterase-5 PET pre-eclampsia

PLGF placental growth factor
PO per os – by mouth
pRBCs packed red blood cells
PrCr protein : creatinine ratio

PRECOG pre-eclampsia community guideline

PRES posterior reversible leukoencephalopathy syndrome

QID four times daily dosing

QLD Queensland Maternity and Neonatal Clinical Guidelines Program

RCT randomised controlled trial RDA recommended daily allowance sFlt-1 soluble fins-like tyrosine kinase-1

sBP systolic blood pressure sEng soluble endoglin SFH symphysis–fundal heigi

SFH symphysis-fundal height SGA small for gestational age sGC soluble guanylate cyclase

SOGC Society of Obstetricians and Gynaecologists of Canada

TID three times daily dosing
TTE transthoracic echocardiography
UNFPA United Nations Population Fund

US United States

VEGF vascular endothelial growth factor WHO World Health Organization